

Axioms for the Set \mathbb{R} of Real Numbers

Axiom 1 (AE: Addition exists). If $a, b \in \mathbb{R}$, then the sum of a and b , denoted by $a + b$, is a uniquely defined number in \mathbb{R} .

Axiom 2 (AA: Addition is associative). For all $a, b, c \in \mathbb{R}$ we have $a + (b + c) = (a + b) + c$.

Axiom 3 (AC: Addition is commutative). For all $a, b \in \mathbb{R}$ we have $a + b = b + a$.

Axiom 4 (AZ: Addition has 0). There is an element 0 in \mathbb{R} such that $0 + a = a + 0 = a$ for all $a \in \mathbb{R}$.

Axiom 5 (AO: Addition has opposites). If $a \in \mathbb{R}$, then the equation $a + x = 0$ has a solution $-a \in \mathbb{R}$. The number $-a$ is called the *opposite* of a .

Axiom 6 (ME: Multiplication exists). If $a, b \in \mathbb{R}$, then the product of a and b , denoted by ab , is a uniquely defined number in \mathbb{R} .

Axiom 7 (MA: Multiplication is associative). For all $a, b, c \in \mathbb{R}$ we have $a(bc) = (ab)c$.

Axiom 8 (MC: Multiplication is commutative). For all $a, b \in \mathbb{R}$ we have $ab = ba$.

Axiom 9 (MO: Multiplication has 1). There is an element $1 \neq 0$ in \mathbb{R} such that $1 \cdot a = a \cdot 1 = a$ for all $a \in \mathbb{R}$.

Axiom 10 (MR: Multiplication has reciprocals). If $a \in \mathbb{R}$ is such that $a \neq 0$, then the equation $a \cdot x = 1$ has a solution $a^{-1} = \frac{1}{a}$ in \mathbb{R} . The number $a^{-1} = \frac{1}{a}$ is called the *reciprocal* of a .

Axiom 11 (DL: Distributive law, the connection between addition and multiplication). For all $a, b, c \in \mathbb{R}$ we have $a(b + c) = ab + ac$.

Axiom 12 (OE: Order exists). Given any $a, b \in \mathbb{R}$, exactly one of these statements is true: $a < b$, $a = b$, or $b < a$. (The symbol $a \leq b$ stands for $a < b$ or $a = b$.)

Axiom 13 (OT: Order is transitive). Given any $a, b, c \in \mathbb{R}$, if $a < b$ and $b < c$, then $a < c$.

Axiom 14 (OA: Order respects addition). Given any $a, b, c \in \mathbb{R}$, if $a < b$ then $a + c < b + c$.

Axiom 15 (OM: Order respects multiplication). Given any $a, b, c \in \mathbb{R}$, if $a < b$ and $0 < c$, then $ac < bc$.

Axiom 16 (CA: Completeness Axiom). If A and B are nonempty subsets of \mathbb{R} such that for every $a \in A$ and for every $b \in B$ we have $a \leq b$, then there exists $c \in \mathbb{R}$ such that $a \leq c \leq b$ for all $a \in A$ and all $b \in B$.

The end of axioms.
